arXiv:0908.3510v2 [math.RT] 2 Mar 2010

n-REPRESENTATION-FINITE ALGEBRAS AND TWISTED FRACTIONALLY CALABI-YAU ALGEBRAS

MARTIN HERSCHEND AND OSAMU IYAMA

Abstract. In this paper, we study n-representation-finite algebras from the viewpoint of the fractionally Calabi-Yau property. We shall show that all n-representation-finite algebras are twisted fractionally Calabi-Yau. We also show that for any $\ell > 0$, twisted $\frac{n(\ell-1)}{\ell}$-Calabi-Yau algebras of global dimension at most n are n-representation-finite. As an application, we give a construction of n-representation-finite algebras using the tensor product.

The Calabi-Yau (CY) property of triangulated categories was introduced by Kontsevich [Ko] (see also [Ke1]). It has played important roles in the representation theory of algebras, especially in the categorification program of Fomin-Zelevinsky cluster algebras by cluster tilting theory (e.g. [Am, BIRS, BMRRT, GLS1, GLS2, IR, Ke2, Ke3, KR]). The derived categories of finite dimensional non-semisimple algebras are never CY, but they are often *fractionally* CY. Some recent meetings [Ba, Bi, P, T] as well as recent results including [D, DL, IIKNS, IKM, KST, Ke2, Ladk, Le, Mi, MY, Y] suggest that the fractionally CY property becomes more and more important in representation theory, singularity theory, commutative and non-commutative algebraic geometry. The aim of this short paper is to apply the fractionally CY property in the study of n-representation-finite algebras defined below.

There is a natural generalization of the classical notion of representation-finiteness from the viewpoint of a higher analogue of Auslander-Reiten theory [Au, I1, I3] studied by several authors [EH, GLS1, IO1, IO2, Lada, HZ1, HZ2, HZ3].

Definition 0.1. We say that a finite dimensional algebra Λ over a field K is *n-representation-finite* (for a positive integer n) if $\operatorname{gl.dim}\Lambda \le n$ and there exists an *n-cluster tilting Λ-module* M, i.e.

$$\begin{aligned} \operatorname{add} M &= \{X \in \operatorname{mod}\Lambda \mid \operatorname{Ext}^i_\Lambda(M,X)=0 \ \text{ for any } \ 0<i<n\}, \\ &= \{X \in \operatorname{mod}\Lambda \mid \operatorname{Ext}^i_\Lambda(X,M)=0 \ \text{ for any } \ 0<i<n\}. \end{aligned}$$

Clearly 1-cluster tilting Λ-modules are additive generators of $\operatorname{mod}\Lambda$, and so Λ is 1-representation-finite if and only if it is representation-finite and hereditary. The following result [I4, 1.3(b)] plays an important role in this paper.

Proposition 0.2. *Let Λ be an n-representation-finite algebra. Let*

$$\begin{aligned} \tau_n &:= \operatorname{Tor}^\Lambda_n(D\Lambda,-) \simeq D\operatorname{Ext}^n_\Lambda(-,\Lambda) &:& \ \operatorname{mod}\Lambda \to \operatorname{mod}\Lambda, \\ \tau_n^- &:= D\operatorname{Tor}^\Lambda_n(D-,D\Lambda) \simeq \operatorname{Ext}^n_\Lambda(D\Lambda,-) &:& \ mod\Lambda \to \operatorname{mod}\Lambda. \end{aligned}$$

Let $P_1,\dots,P_a$ be the isoclasses of indecomposable projective Λ-modules, and let I_i be the indecomposable injective Λ-module corresponding to P_i.

(a) *There exists a permutation $\sigma \in \mathfrak{S}_a$ and positive integers $\ell_1,\dots,\ell_a$ such that $\tau_n^{\ell_i-1} I_i \simeq P_{\sigma(i)}$ for any i.*

(b) *There exists a unique basic n-cluster tilting Λ-module M, which is given as the direct sum of the following mutually non-isomorphic indecomposable Λ-modules.*

$$\begin{array}{cccccc} I_1, & \tau_n I_1, & \tau_n^2 I_1, & \cdots & \tau_n^{\ell_1-2} I_1, & \tau_n^{\ell_1-1} I_1 \simeq P_{\sigma(1)} \\ I_2, & \tau_n I_2, & \tau_n^2 I_2, & \cdots & \tau_n^{\ell_2-2} I_2, & \tau_n^{\ell_2-1} I_2 \simeq P_{\sigma(2)} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ I_a, & \tau_n I_a, & \tau_n^2 I_a, & \cdots & \tau_n^{\ell_a-2} I_a, & \tau_n^{\ell_a-1} I_a \simeq P_{\sigma(a)} \end{array}$$

(c) *We have mutually quasi-inverse equivalences* $\tau_n : \mathrm{add}(M/\Lambda) \simeq \mathrm{add}(M/D\Lambda)$ *and* $\tau_n^- : \mathrm{add}(M/D\Lambda) \simeq \mathrm{add}(M/\Lambda)$.

Throughout this paper, let K be a field and Λ a finite dimensional K-algebra of finite global dimension. All modules are finitely generated left modules unless stated otherwise. The composition fg of morphisms (or arrows) means that f is first and g is next. We denote by $\mathcal{D} = \mathcal{D}^{\mathrm{b}}(\mathrm{mod}\,\Lambda)$ the bounded derived category of the category $\mathrm{mod}\,\Lambda$ of all Λ-modules. Then the Nakayama functor

$$\nu = \nu_\Lambda := (D\Lambda) \overset{\mathbf{L}}{\otimes}_\Lambda - \simeq D \circ \mathbf{R}\mathrm{Hom}_\Lambda(-, \Lambda) : \mathcal{D} \to \mathcal{D}$$

gives a Serre functor, i.e. there exists a functorial isomorphism

$$\mathrm{Hom}_{\mathcal{D}}(X, Y) \simeq D\,\mathrm{Hom}_{\mathcal{D}}(Y, \nu\, X)$$

for any $X, Y \in \mathcal{D}$ [H, BK]. For each K-algebra endomorphism ϕ of Λ, we define an endofunctor of $\mathcal{D}$ by

$$\phi^* := \Lambda_\phi \overset{\mathbf{L}}{\otimes}_\Lambda - : \mathcal{D} \to \mathcal{D},$$

where we denote by Λ_ϕ the $\Lambda \otimes_K \Lambda^{\mathrm{op}}$-module such that the right action is given by $a \cdot b := a\phi(b)$. For any $x \in \mathrm{Ker}\,\phi$, the morphism $\Lambda \to \Lambda$, $\lambda \mapsto \lambda x$ is mapped to 0 by ϕ^*. Thus ϕ^* is an autofunctor if and only if ϕ is an automorphism.

Definition 0.3. We say that Λ is *twisted fractionally CY* (or *twisted* $\frac{m}{\ell}$*-CY*) if there exists an isomorphism

$$\nu^\ell \simeq [m] \circ \phi^* \tag{1}$$

of functors for some integers $\ell \neq 0$ and m and a K-algebra endomorphism ϕ of Λ. Then ϕ must be an automorphism since ν is an autofunctor. When $\phi = \mathrm{id}$, we say that Λ is *fractionally CY* (or $\frac{m}{\ell}$*-CY*).

Notice that (twisted) $\frac{m}{\ell}$-CY algebras are also (twisted) $\frac{mk}{\ell k}$-CY for any $k \neq 0$. The converse does not hold in general, but the Calabi-Yau dimension is unique as a rational number.

Acknowledgement The authors would like to thank an anonymous referee for careful reading the first draft and helpful comments. The second author was supported by JSPS Grant-in-Aid for Scientific Research 21740010 and 60015849. The first author is grateful to JSPS for funding his stay at Nagoya University, during which this paper was written.

1. MAIN RESULTS

Our first result is the following.

Theorem 1.1. *Let* Λ *be a ring-indecomposable* n*-representation-finite algebra. Then:*

(a) Λ *is twisted fractionally CY.*

(b) *More precisely, let* a *be the number of simple* Λ*-modules, and let* b *be the number of idecomposable summands of the basic* n*-cluster tilting* Λ*-module. Then* Λ *is twisted* $\frac{n(b-a)k}{bk}$*-CY for some* $k > 0$.

Our second result shows that a certain converse of the statement (a) above holds.

Definition 1.2. We say that an n-representation-finite algebra Λ is ℓ*-homogeneous* (or simply *homogeneous*) if $\ell := \ell_1 = \cdots = \ell_a$ holds in the notation of Proposition 0.2. In this case, we have $\frac{b}{a} = \ell$ for a and b in Theorem 1.1(b).

Theorem 1.3. *For a finite dimensional algebra* Λ *and a positive integer* ℓ*, the following conditions are equivalent.*

(a) Λ *is* ℓ*-homogeneous* n*-representation-finite.*

(b) Λ *is twisted* $\frac{n(\ell-1)}{\ell}$*-CY and* $\mathrm{gl.dim}\,\Lambda \leq n$.

A typical example is given by tubular algebras of type $(2, 2, 2, 2)$ [R] for the case $n = \ell = 2$. They are known to be $\frac{2}{2}$-CY [LM], and also to be 2-representation-finite [IO2].

Let us apply these results to construct n-representation-finite algebras using the tensor product. The next observation shows that fractionally CY algebras behave nicely under the tensor product, where we denote $\otimes_K$ by $\otimes$. The requirement on the field K to be perfect is needed to guarantee that the tensor product of two finite dimensional K-algebras of finite global dimension is still of finite global dimension.

Proposition 1.4. *Let K be a perfect field. If Λ_i is $\frac{m_i}{\ell_i}$-CY (respectively, twisted $\frac{m_i}{\ell_i}$-CY), then $\Lambda_1 \otimes \cdots \otimes \Lambda_k$ is $\frac{m}{\ell}$-CY (respectively, twisted $\frac{m}{\ell}$-CY) for the least common multiple ℓ of $\ell_1, \ldots, \ell_k$ and $m := \ell(\frac{m_1}{\ell_1} + \cdots + \frac{m_k}{\ell_k})$.*

As an easy consequence, we have the following useful result.

Corollary 1.5. *Let K be a perfect field and ℓ a positive integer. If Λ_i is ℓ-homogeneous n_i-representation-finite, then $\Lambda_1 \otimes \cdots \otimes \Lambda_k$ is an ℓ-homogeneous $(n_1 + \cdots + n_k)$-representation-finite algebra with an $(n_1 + \cdots + n_k)$-cluster tilting module $\bigoplus_{i=0}^{\ell-1}(\tau_{n_1}^i \Lambda_1 \otimes \cdots \otimes \tau_{n_k}^i \Lambda_k)$.*

Notice that the statement is not true if we drop the assmption of ℓ-homogeneity (see Remark 1.6(a)).

The relationship between n-representation-finiteness and the factionally CY property is summarized in the following diagram:


ℓ-homogeneous n-representation-finite
Thm.1.3
twisted $\frac{n(\ell-1)}{\ell}$-CY of global dimension at most n
n-representation-finite
Thm.1.1
twisted fractionally CY
?Rem.1.6(b)
×Rem.1.6(a)
?Rem.1.6(b)
fractionally CY

Remark 1.6. (a) Let $\Lambda := K[\bullet \to \bullet] \otimes K[\bullet \to \bullet]$. Then Λ is fractionally CY, but not n-representation-finite for any n. Moreover Λ is derived equivalent to the path algebras of type D_4, which are 1-representation-finite. In particular we have the following conclusions.
  - There exist fractionally CY algebras which are not n-representation-finite for any n.
  - n-representation-finiteness is not preserved under derived equivalence.
  - n-representation-finiteness is not preserved under the tensor product.

(b) Although we already know that every n-representation-finite algebra is twisted fractionally CY, we do not know the answer to the following question.
  - Is every n-representation-finite algebra fractionally CY?

Also we do not know the answer to the following related question.
  - Is every twisted fractionally CY algebra fractionally CY?

This is equivalent to that ϕ in (1) has a finite order in the outer automorphism group of Λ.

(c) Since the Serre functor is unique up to isomorphism, the fractionally CY property is invariant under derived equivalence. However, we do not know the answer to the following question.
  - Are twisted fractionally CY algebras closed under derived equvalence?

If we fix the CY dimension, then the answer is negative. For example $K[\bullet \to \bullet \leftarrow \bullet]$ is twisted $\frac{1}{2}$-CY, but $K[\bullet \to \bullet \to \bullet]$ is not (see section 3.1).

We end this section by an observation on endomorphism algebras $\Gamma := \mathrm{End}_\Lambda(M)$ of n-cluster tilting Λ-modules M. They are called *n-Auslander algebras*, and satisfies $\mathrm{gl.dim}\,\Gamma \leq n+1 \leq \mathrm{dom.dim}\,\Gamma$ [I2]. We have the following descrption of Γ when Λ is ℓ-homogeneous n-representation-finite.

Proposition 1.7. *If Λ in an ℓ-homogeneous n-representation-finite algebra, then the n-Auslander algebra of Λ is described as*

$$\begin{pmatrix} \Lambda & T & \cdots & T^{\otimes_\Lambda(\ell-2)} & T^{\otimes_\Lambda(\ell-1)} \\ 0 & \Lambda & \cdots & T^{\otimes_\Lambda(\ell-3)} & T^{\otimes_\Lambda(\ell-2)} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & \Lambda & T \\ 0 & 0 & \cdots & 0 & \Lambda \end{pmatrix}$$

for the $\Lambda \otimes \Lambda^{\mathrm{op}}$-module $T := \mathrm{Ext}^n_\Lambda(D\Lambda, \Lambda)$.

By Corollary 1.5, the $(n_1+\cdots+n_k)$-Auslander algebra of $\Lambda_1 \otimes \cdots \otimes \Lambda_k$ is given by the 'component-wise tensor product'

$$\begin{pmatrix} \Lambda_1 \otimes \cdots \otimes \Lambda_k & T_1 \otimes \cdots \otimes T_k & \cdots & T_1^{\otimes_{\Lambda_1}(\ell-2)} \otimes \cdots \otimes T_k^{\otimes_{\Lambda_k}(\ell-2)} & T_1^{\otimes_{\Lambda_1}(\ell-1)} \otimes \cdots \otimes T_k^{\otimes_{\Lambda_k}(\ell-1)} \\ 0 & \Lambda_1 \otimes \cdots \otimes \Lambda_k & \cdots & T_1^{\otimes_{\Lambda_1}(\ell-3)} \otimes \cdots \otimes T_k^{\otimes_{\Lambda_k}(\ell-3)} & T_1^{\otimes_{\Lambda_1}(\ell-2)} \otimes \cdots \otimes T_k^{\otimes_{\Lambda_k}(\ell-2)} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & \Lambda_1 \otimes \cdots \otimes \Lambda_k & T_1 \otimes \cdots \otimes T_k \\ 0 & 0 & \cdots & 0 & \Lambda_1 \otimes \cdots \otimes \Lambda_k \end{pmatrix}$$

of the n_i-Auslander algebras. This is interesting since in general the tensor product of higher Auslander algebras is not a higher Auslander algebra again.

2. HOMOGENEITY AND $(n+1)$-PREPROJECTIVE ALGEBRAS

In this section, we further study the homogeneity of n-representation-finite algebras.

The following result gives a useful criterion for n-representation-finite algebras to be homogeneous.

Proposition 2.1. *Let Λ be a ring-indecomposable n-representation-finite algebra. Then Λ is homogeneous if and only if $\ell_i = \ell_{\sigma(i)}$ holds for any i.*

We put

$$\nu_n := \nu \circ [-n] : \mathcal{D} \to \mathcal{D}.$$

For an arbitrary n-representation-finite algebra Λ, we denote by

$$\Pi := \bigoplus_{i \in \mathbf{Z}} \mathrm{Hom}_{\mathcal{D}}(\Lambda, \nu_n^{-i} \Lambda)$$

the corresponding $(n+1)$*-preprojective algebra*. This is a positively graded K-algebra, and the multiplication of $f \in \mathrm{Hom}_{\mathcal{D}}(\Lambda, \nu_n^{-i} \Lambda)$ and $g \in \mathrm{Hom}_{\mathcal{D}}(\Lambda, \nu_n^{-j} \Lambda)$ is given by

$$f \cdot g := f\, \nu_n^{-i}\, g.$$

Then Π can be viewed as the tensor algebra over Λ of the bimodule $\mathrm{Ext}^n_\Lambda(D\Lambda, \Lambda)$ [IO1, 2.12]. Notice that Π is the cohomology in degree zero of the DG algebra defined in [Ke3]. It is shown in [IO2] that Π is a finite dimensional selfinjective algebra.

A *Nakayama automorphism* ψ of Π is an automorphism which gives an isomorphism $D\Pi \simeq \Pi_\psi$ of $\Pi \otimes \Pi^{\mathrm{op}}$-modules. This is uniquely determined as an element of the outer automorphism group of Π. Moreover

$$\psi^* = \Pi_\psi \otimes_\Pi - \simeq (D\Pi) \otimes_\Pi - \simeq D\,\mathrm{Hom}_\Pi(-, \Pi) : \mathrm{mod}\,\Pi \to \mathrm{mod}\,\Pi$$

gives the Nakayama functor.

The next result gives the relationship between ψ and σ. Notice that the isoclasses of indecomposable projective Π-modules are give by $\Pi \otimes_\Lambda P_i$ for $1 \leq i \leq a$.

Proposition 2.2. *Let ψ be a Nakayama automorphism of Π, and let σ be as in Proposition 0.2. Then we have $\psi^*(\Pi \otimes_\Lambda P_i) \simeq \Pi \otimes_\Lambda P_{\sigma(i)}$ for any i. In particular σ gives the Nakayama permutation of Π.*

The following main result in this section characterizes homogeneity of n-representation-finite algebras in terms of the corresponding $(n+1)$-preprojective algebras.

Theorem 2.3. *Let ψ be a Nakayama automorphism of Π, and let I be the ideal of Π such that $\Lambda = \Pi/I$. Then Λ is homogeneous if and only if $\psi(I) = I$.*

When Λ is homogeneous, we can interpret a Nakayama automorphism of Π as follows: By Theorem 1.3, there exists a K-algebra automorphism ϕ of Λ such that $\nu^\ell \simeq [n(\ell-1)] \circ \phi^*$ for some $\ell \geq 1$.

Proposition 2.4. *Assume that Λ is homogeneous. Then the automorphism ϕ of Λ extends to a graded K-algebra automorphism $\widetilde{\phi}$ of Π which is a Nakayama automorphism.*

3. Examples

3.1. **Path algebras.** Let us consider Dynkin diagrams:


A_n $1 — 2 — 3 —$ …… $— n-2 — n-1 — n$
D_n $1 — 2 — 3 —$ …… $— n-2 — n-1$, n
E_6 $1 — 2 — 3 — 5 — 6$, 4
E_7 $1 — 2 — 3 — 4 — 5 — 6$, 7
E_8 $1 — 2 — 3 — 4 — 5 — 6 — 7$, 8

The Coxeter number h of each Dynkin diagram is given as follows:

A_n	D_n	E_6	E_7	E_8
$n+1$	$2(n-1)$	12	18	30

The following proposition is well-known [MY, 4.1].

Proposition 3.1. *Let Q be an acyclic quiver. Then KQ is (twisted) fractionally CY if and only if Q is a Dynkin quiver. In this case, KQ is $\frac{h-2}{h}$-CY, and if Q is D_n with even n, E_7 or E_8, then KQ is $\frac{\frac{h}{2}-1}{\frac{h}{2}}$-CY.*

We define an involution ω of each Dynkin diagram as follows:

- For A_n, we put $\omega(i) = n+1-i$.
- For D_n with odd n, we put $\omega(n-1) = n$, $\omega(n) = n-1$ and $\omega(i) = i$ for other i.
- For E_6, we put $\omega(1) = 6$, $\omega(2) = 5$, $\omega(5) = 2$, $\omega(6) = 1$ and $\omega(i) = i$ for other i.
- For other types, we put $\omega = \mathrm{id}$.

Proposition 3.2. *Let Q be an acyclic quiver.*

(a) *The following conditions are equivalent.*
  (i) *KQ is ℓ-homogeneous 1-representation-finite for some ℓ.*
  (ii) *Q is a Dynkin quiver and the orientation is stable under ω.*
(b) *If the conditions in (a) are satisfied, then $\ell = \frac{h}{2}$ and KQ is twisted $\frac{\frac{h}{2}-1}{\frac{h}{2}}$-CY and the twist is induced by ω. In particular, if $\omega = \mathrm{id}$, then KQ is $\frac{\frac{h}{2}-1}{\frac{h}{2}}$-CY.*
(c) *The underlying diagrams of ω-stable Dynkin quivers are classified by ℓ as follows.*

ℓ	$\ell\ (\neq 6,9,15)$	6	9	15
Q	$A_{2\ell-1}, D_{\ell+1}$	A_{11}, D_7, E_6	A_{17}, D_{10}, E_7	A_{29}, D_{16}, E_8

Proof. (a) This follows from Gabriel's description of Auslander-Reiten quivers of KQ [G].

(b) The equality $\ell = \frac{h}{2}$ is well-known [G], and the latter assertion follows from Theorem 1.3. ☐

By Corollary 1.5, for any choice of homogeneous algebras $\Lambda_1, \dots, \Lambda_k$ which belong to the same column in the table in (c) above, we have an ℓ-homogeneous k-representation-finite algebra $\Lambda_1 \otimes \cdots \otimes \Lambda_k$.

As one of the simplest examples, we have the following 2-representation-finite algebras.



3.2. **n-representation-finite algebras of type A.** We give a different class of examples of ℓ-homogeneous n-representation-finite algebras based on results in [IO1]. Fix positive integers n and s. We define the quiver $Q := Q^{(n,s)}$ with the set Q_0 of vertices and the set Q_1 of arrows by

$$\begin{aligned} Q_0 &:= \{x = (x_1, x_2, \dots, x_{n+1}) \in \mathbf{Z}^{n+1}_{\geq 0} \mid \sum_{i=1}^{n+1} x_i = s - 1\}, \\ Q_1 &:= \{x \xrightarrow{i} x + f_i \mid 1 \leq i \leq n+1,\ x, x + f_i \in Q_0\} \end{aligned}$$

where f_i denotes the vector $f_i := (0, \dots, 0, \overset{i}{-1}, \overset{i+1}{1}, 0, \dots, 0)$ for $1 \leq i \leq n$ and $f_{n+1} := (\overset{1}{1}, 0, \dots, 0, \overset{n+1}{-1})$.

Example 3.3. $Q^{(1,5)}$, $Q^{(2,4)}$ and $Q^{(3,3)}$ are the following quivers.



We define the K-algebra $\Gamma = \Gamma^{(n,s)} := KQ/I$, where I is the ideal defined by the following relations: For any $x \in Q_0$ and $1 \leq i, j \leq n+1$ satisfying $x + f_i, x + f_i + f_j \in Q_0$,

$$(x \xrightarrow{i} x + f_i \xrightarrow{j} x + f_i + f_j) = \begin{cases} (x \xrightarrow{j} x + f_j \xrightarrow{i} x + f_i + f_j) & \text{if } x + f_j \in Q_0, \\ 0 & \text{otherwise.} \end{cases}$$

We call a subset C of Q_1 a *cut* if it contains exactly one arrow from each cycle of length $n+1$ in Q. In this case, we define a K-algebra by

$$\Lambda_C := \Gamma / \langle C \rangle,$$

for the ideal $\langle C \rangle$ of Γ generated by the arrows in C. A main result in [IO1] is the following. (Notice that the assumption in [IO1] that K is algebraically closed is not necessary.)

Proposition 3.4. *For any cut C of Q, the algebra Λ_C is an n-representation-finite algebra with the $(n+1)$-preprojective algebra Γ.*

It is natural to ask when Λ_C is homogeneous. To answer this question we introduce the automorphism ω of Q defined by

$$\omega(x_1, x_2, \dots, x_{n+1}) := (x_{n+1}, x_1, \dots, x_n).$$

We leave the proof of the following statement to the reader.

Theorem 3.5. *The quiver morphism ω induces a Nakayama automorphism of Γ.*

Proof. For each $x = (x_1, x_2, \dots, x_{n+1}) \in \mathbf{Z}^{n+1}$ we adopt the convention $x_{i+n+1} = x_i$. Now let $x \in Q_0$ and $1 \leq i \leq n+1$ such that $x + f_i \in Q_0$. Then $\omega(x + f_i) = \omega(x) + f_{i+1} \in Q_0$ and

$$\omega(x \xrightarrow{i} x + f_i) = \omega(x) \xrightarrow{i+1} \omega(x) + f_{i+1}.$$

Now assume $x + f_i + f_j \in Q_0$. Then

$$\omega(x \xrightarrow{i} x + f_i \xrightarrow{j} x + f_i + f_j) = \omega(x) \xrightarrow{i+1} \omega(x) + f_{i+1} \xrightarrow{j+1} \omega(x) + f_{i+1} + f_{j+1}.$$

Since $x + f_j \in Q_0$ if and only if $\omega(x) + f_{j+1} \in Q_0$ it follow that ω induces an automorphism on Γ.

The relations defining I are only commutativity and zero relations. Thus the set B of non-zero residue classes $p + I \neq I$ of paths p in Q forms a basis of Γ. Since Γ is finite dimensional, $D\Gamma$ has basis $DB := \{\bar{b}^* | \bar{b} \in B\}$ dual to B. We proceed to show that there is a bijection

$$\phi : DB \simeq B$$

that extends to a bimodule isomorphism $D\Gamma \simeq \Gamma_\omega$.

Define the quiver $\widehat{Q}$ by

$$\begin{aligned} \widehat{Q}_0 &:= \{x \in \mathbf{Z}^{n+1} \mid \sum_{i=1}^{n+1} x_i = s - 1\}, \\ \widehat{Q}_1 &:= \{x \xrightarrow{i} x + f_i \mid 1 \leq i \leq n + 1\}. \end{aligned}$$

Hence Q is the full subquiver of $\widehat{Q}$ with vertecies $\{x \in \widehat{Q}_0 \mid x_i \geq 0 \text{ for all } 1 \leq i \leq n+1\}$. Let $\widehat{I}$ be the ideal in $K\widehat{Q}$ defined by the relations

$$(x \xrightarrow{i} x + f_i \xrightarrow{j} x + f_i + f_j) = (x \xrightarrow{j} x + f_j \xrightarrow{i} x + f_i + f_j)$$

and set $\widehat{\Gamma} = K\widehat{Q}/\widehat{I}$. We have a canonical surjective ring morphism $\pi : \widehat{\Gamma} \to \Gamma$ with kernel

$$K = \sum_{z \notin Q_0} \widehat{\Gamma} e_z \widehat{\Gamma},$$

where e_z denotes the path of length 0 at z. Since $\widehat{I}$ is defined by commutativity relations, the residue classes of paths form a basis of $\widehat{\Gamma}$. Moreover, the residue classes of paths that are not in K are mapped bijectively to B by π.

We define a $\mathbf{Z}^{n+1}$-grading g on $\widehat{\Gamma}$ by $(g(x \xrightarrow{i} x + f_i))_j = \delta_{ij}$. It is well-defined since the defining relations of $\widehat{I}$ are homogeneous. For paths p, p' in $\widehat{Q}$ we write $p \equiv p'$ if and only if $p - p' \in \widehat{I}$. Let p be a path from x to y of degree d. Then $y - x = \sum_i d_i f_i$. In fact $p \equiv p_{x,d,y}$, where

$$p_{x,d,y} = x \xrightarrow{1} \cdots \xrightarrow{1} x + d_1 f_1 \xrightarrow{2} \cdots \xrightarrow{2} x + d_1 f_1 + d_2 f_2 \xrightarrow{3} \quad \cdots \quad \xrightarrow{n} y - d_{n+1} f_{f+1} \xrightarrow{n+1} \cdots \xrightarrow{n+1} y.$$

In particular, $p + \widehat{I}$ is determined by d together with either x or y. Moreover, for each path p' from x to z, there is a path q from z to y such that $p \equiv p'q$ if and only if $g(p')_i \leq d_i$ for all $1 \leq i \leq n + 1$ (take $q = p_{z,d-g(p'),y}$).

Observe that $p + \widehat{I} \in K$ if and only if there are paths p' from x to z and q from z to y such that $p \equiv p'q$ and $z \notin Q_0$. By the above observation this is equivalent to that there is $a \in \mathbf{Z}^{n+1}_{\geq 0}$ such that $a_j \leq d_j$ for each $1 \leq j \leq n + 1$ and $x + \sum_i a_i f_i \notin Q_0$. This holds if and only if there is $1 \leq j \leq n + 1$ such that $x_j < d_j$ (take $a_i = \delta_{ij} d_j$).

We conclude that for every path p in Q from x to z such that $p + I \neq I$ there is a path q in Q from z to some vertex y such that $g(pq) = x$. In particular, $pq + I \neq I$ and

$$y = x + \sum_i x_i f_i.$$

Thus

$$y_j = x_j - x_j + x_{j-1} = x_{j-1} = \omega(x)_j.$$

Moreover, $q + I$ is determined by p since $g(q) = x - g(p)$. Similarly, for each path q in Q from z to $\omega(x)$ such that $q \notin I$ there is a path p in Q from x to z, unique up to $\equiv$ such that $g(pq) = x$. We define the bijection $\phi : DB \simeq B$ by $\phi((p + I)^*) = q + I$, where p is a path from x to z and q is a path from z such that $g(pq) = x$. It induces a linear bijection

$$\phi : D\Gamma \simeq \Gamma_\omega.$$

We proceed to show that ϕ is a bimodule isomorphism, thus completing the proof.

First consider the following configuration of paths in Q.

$$\begin{tikzcd} x \arrow[r, dashed, "a"] & y \arrow[r, dashed, "p'"] & z \arrow[r, dashed, "b"] & w \arrow[r, dashed, "q"] & \omega(x) \arrow[r, dashed, "\omega(a)"] & \omega(y) \end{tikzcd}$$

Then

$$g(p'bq\omega(a)) - g(ap'bq) = g(\omega(a)) - g(a) = \sum_i g(a)_i f_i = y - x.$$

In particular, $g(ap'bq) = x$ if and only if $g(p'bq\omega(a)) = y$, or equivalently $\phi((ap'b+I)^*) = q+I$ if and only if $\phi((p'+I)^*) = bq\omega(a)+I$.

Now fix paths α, β, p from x to w and q from w to $\omega(x)$ such that $g(pq) = x$. Set $\bar{b} = (p+I)$ so that $\phi(\bar{b}^*) = q+I$. We claim that there is p' such that $p \equiv \alpha p'\beta$ if and only if $\beta q\omega(\alpha) \notin I$. If such p' exists then $g(\alpha p'\beta q) = x$ and by our observation $g(p'\beta q\omega(\alpha)) = y$. This implies $\beta q\omega(\alpha) \notin I$. On the other hand, if $\beta q\omega(\alpha) \notin I$, then there is p' such that $g(p'\beta q\omega(\alpha)) = y$ and $g(\alpha p'\beta q) = x$ which implies $\alpha p'\beta \equiv p$.

If there is a p' satisfying $p \equiv \alpha p'\beta$, then by the bimodule structure on $D\Gamma$, we have $\beta\bar{b}^*\alpha = (p'+I)^*$. Otherwise $\beta\bar{b}^*\alpha = 0$. In either case $\phi(\beta\bar{b}^*\alpha) = \beta q\omega(\alpha)+I = \beta\phi(\bar{b}^*)\alpha$ by the above observations. □

We have the following characterization.

Theorem 3.6. *Let C be an admissible set of Q. Then the n-representation-finite algebra Λ_C is homogeneous if and only if $\omega(C) = C$. In this case, we have $\ell = \frac{s-1}{n+1} + 1$.*

Proof. The first assertion follows from Theorems 2.3 and 3.5. To show the second assertion we apply Theorem 1.1(a). We know that a is equal to the number $\binom{s+n-1}{n}$ of vertices of the quiver $Q^{(n,s)}$, and b is equal to the number $\binom{s+n}{n+1}$ of vertices of the quiver $Q^{(n+1,s)}$ by [I4, 6.12]. Thus $\ell = \frac{b}{a} = \frac{s+n}{n+1}$. □

For the case $n = 1$, this result gives the characterization of type A_s quivers given in Proposition 3.2.

For the case $n = 2$ and $s = 4$, there are the following five 2-homogeneous 2-representation-finite algebras



4. Proof of our results

Recall that Λ is a finite dimensonal K-algebra of finite global dimension and $\mathcal{D} = \mathcal{D}^{\mathrm{b}}(\operatorname{mod}\Lambda)$ is the bounded derived category of $\operatorname{mod}\Lambda$.

We need the following observation.

Lemma 4.1. *For a $\Lambda\otimes\Lambda^{\mathrm{op}}$-module X, the following conditions are equivalent.*

(a) *$X \simeq \Lambda$ as Λ-modules.*

(b) *$X \simeq \Lambda_\phi$ as $\Lambda\otimes\Lambda^{\mathrm{op}}$-modules for some K-algebra endomorphism ϕ of Λ.*

Proof. (b)⇒(a) This is clear.

(a)⇒(b) Fix an isomorphism $f \in \operatorname{Hom}_\Lambda(X,\Lambda)$. For each $\lambda \in \Lambda$, the right multiplication $(\cdot\lambda) \in \operatorname{End}_\Lambda(X)$ gives an endomorphism $f^{-1}\cdot(\cdot\lambda)\cdot f \in \operatorname{End}_\Lambda(\Lambda)$. This is given by the right multiplication of an element $\phi(\lambda) \in \Lambda$. Then ϕ gives the desired K-algebra endomorphism of Λ. □

We need the following observation.

Lemma 4.2. *Let $X \in \mathcal{D}^{\mathrm{b}}(\operatorname{mod}\Lambda\otimes\Lambda^{\mathrm{op}})$.*

(a) *The following conditions are equivalent.*

  (i) *$X \simeq \Lambda$ in $\mathcal{D}^{\mathrm{b}}(\operatorname{mod}\Lambda)$.*

  (ii) *$X \simeq \Lambda_\phi$ in $\mathcal{D}^{\mathrm{b}}(\operatorname{mod}\Lambda\otimes\Lambda^{\mathrm{op}})$ for some K-algebra endomorphism ϕ of Λ.*

(b) *The following conditions are equivalent.*
  (i) *There exists an isomorphism* $f : \Lambda \to X$ *in* $\mathcal{D}^{\mathrm{b}}(\operatorname{mod}\Lambda)$ *which commutes with the right action of* Λ.
  (ii) $X \simeq \Lambda$ *in* $\mathcal{D}^{\mathrm{b}}(\operatorname{mod}\Lambda \otimes \Lambda^{\mathrm{op}})$.

Proof. (a) The condition (i) is equivalent to the following one:

- $H^i(X) = 0$ for any $i \neq 0$ and $H^0(X) \simeq \Lambda$ as Λ-modules.

In view of Lemma 4.1, this is equivalent to the following condition:

- $H^i(X) = 0$ for any $i \neq 0$ and $H^0(X) \simeq \Lambda_\phi$ as $\Lambda \otimes \Lambda^{\mathrm{op}}$-modules for some K-algebra endomorphism ϕ of Λ.

This is equivalent to the condition (ii).

(b) The condition (i) is equivalent to the following one:

- $H^i(X) = 0$ for any $i \neq 0$ and $H^0(X) \simeq \Lambda$ as $\Lambda \otimes \Lambda^{\mathrm{op}}$-modules.

This is equivalent to the condition (ii). □

Let us show the following characterization of (twisted) fractionally CY property.

Proposition 4.3. *Let* Λ *be a finite dimensional* K*-algebra of finite global dimension.*

(a) *The following conditions are equivalent.*
  (i) Λ *is twisted* $\frac{m}{\ell}$*-CY.*
  (ii) $\nu^\ell\,\Lambda \simeq \Lambda[m]$ *in* $\mathcal{D}^{\mathrm{b}}(\operatorname{mod}\Lambda)$.
  (iii) *There exists an isomorphism* $(D\Lambda)^{\overset{\mathbf{L}}{\otimes}_\Lambda \ell} \simeq \Lambda[m]$ *in* $\mathcal{D}^{\mathrm{b}}(\operatorname{mod}\Lambda)$.
(b) *The following conditions are equivalent.*
  (i) Λ *is* $\frac{m}{\ell}$*-CY.*
  (ii) $\nu^\ell \simeq [m]$ *as functors on* $\mathcal{D}^{\mathrm{b}}(\operatorname{mod}\Lambda)$.
  (iii) *There exists an isomorphism* $(D\Lambda)^{\overset{\mathbf{L}}{\otimes}_\Lambda \ell} \simeq \Lambda[m]$ *in* $\mathcal{D}^{\mathrm{b}}(\operatorname{mod}\Lambda \otimes \Lambda^{\mathrm{op}})$.

Proof. (a) By definition (ii) is equivalent to (iii). Since $\Lambda_\phi \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda[m] \simeq \Lambda[m]$, (i) implies (ii).

If (iii) holds, then $(D\Lambda)^{\overset{\mathbf{L}}{\otimes}_\Lambda \ell} \simeq \Lambda_\phi[m]$ in $\mathcal{D}^{\mathrm{b}}(\operatorname{mod}\Lambda \otimes \Lambda^{\mathrm{op}})$ for some K-algebra endomorphism ϕ of Λ by Lemma 4.2(a). Thus we have $\nu^\ell = [m] \circ \phi^*$, and (i) holds.

(b) By definition (i) is equivalent to (ii). Since $\nu^\ell = (D\Lambda)^{\overset{\mathbf{L}}{\otimes}_\Lambda \ell} \overset{\mathbf{L}}{\otimes}_\Lambda -$ and $[m] \simeq \Lambda[m] \overset{\mathbf{L}}{\otimes}_\Lambda -$, (iii) implies (ii). Finally, if (ii) holds, then there exists an isomorphism $f : (D\Lambda)^{\overset{\mathbf{L}}{\otimes}_\Lambda \ell} \to \Lambda[m]$ in $\mathcal{D}^{\mathrm{b}}(\operatorname{mod}\Lambda)$ which commutes with the right action of Λ. By Lemma 4.2(b), we have that (iii) holds. □

We need the following observations.

Lemma 4.4. (a) *Let* Λ *be a finite dimensional* K*-algebra with* $\operatorname{gl.dim}\Lambda \leq n$ *and* $X \in \operatorname{mod}\Lambda$. *Then:*

- $\tau_n X \simeq \nu_n X$ *if and only if* $\operatorname{Ext}^i_\Lambda(X, \Lambda) = 0$ *for any* $i \neq n$.
- *If* X *is indecomposable non-projective and satisfies* $\operatorname{Ext}^i_\Lambda(X, \Lambda) = 0$ *for any* $0 < i < n$*, then we have* $\operatorname{Hom}_\Lambda(X, \Lambda) = 0$ *and* $\tau_n X \simeq \nu_n X$.
- $\tau_n^- X \simeq \nu_n^{-1} X$ *if and only if* $\operatorname{Ext}^i_\Lambda(D\Lambda, X) = 0$ *for any* $i \neq n$.
- *If* X *is indecomposable non-injective and satisfies* $\operatorname{Ext}^i_\Lambda(D\Lambda, X) = 0$ *for any* $0 < i < n$*, then we have* $\operatorname{Hom}_\Lambda(D\Lambda, X) = 0$ *and* $\tau_n^- X \simeq \nu_n^{-1} X$.

(b) *Let* Λ *be an* n*-representation-finite algebra and* M *an* n*-cluster tilting* Λ*-module. Then we have*

$$\tau_n X \simeq \nu_n X \quad \text{and} \quad \operatorname{Hom}_\Lambda(X, \Lambda) = 0 \quad (\text{respectively,} \quad \tau_n^- X \simeq \nu_n^{-1} X \quad \text{and} \quad \operatorname{Hom}_\Lambda(D\Lambda, X) = 0)$$

for any $X \in \operatorname{add} M$ *without non-zero projective (respectively, injective) direct summands.*

(c) *Let* Λ *be a finite dimensional* K*-algebra. For any* K*-algebra automorphism* ϕ *of* Λ*, we have an isomorphism* $\phi^*\,\nu \simeq \nu\,\phi^*$ *of autofunctors of* $\mathcal{D}$.

Proof. (a) The first and the third assertions are clear. For the remaining assertions, see [I4, 2.3(b)].

(b) This is immediate from (a) and the definition of n-cluster tilting modules.

(c) There exist isomorphisms

$$\Lambda_\phi \otimes_\Lambda (D\Lambda) \simeq {}_{\phi^{-1}}(D\Lambda) \simeq D(\Lambda_{\phi^{-1}}) \simeq D({}_\phi\Lambda) \simeq (D\Lambda)_\phi \simeq (D\Lambda) \otimes_\Lambda \Lambda_\phi$$

of $\Lambda \otimes \Lambda^{\mathrm{op}}$-modules. This gives an isomorphism $\Lambda_\phi \overset{\mathbf{L}}{\otimes}_\Lambda (D\Lambda) \simeq (D\Lambda) \overset{\mathbf{L}}{\otimes}_\Lambda \Lambda_\phi$ in $\mathcal{D}^{\mathrm{b}}(\operatorname{mod} \Lambda \otimes \Lambda^{\mathrm{op}})$. Thus we have the desired isomorphism. □

4.1. **Proof of Theorem 1.1.** We use the notation in Proposition 0.2. By Lemma 4.4(b), we have

$$\nu_n^{\ell_i - 1} I_i \simeq P_{\sigma(i)}$$

for any $1 \le i \le a$. In particular, we have

$$\nu^{\ell_i} P_i \simeq \nu^{\ell_i - 1} I_i \simeq P_{\sigma(i)}[n(\ell_i - 1)]$$

for any $1 \le i \le a$. We take $c > 0$ such that $\sigma^c = \mathrm{id}$, and let

$$m_i := \ell_i + \ell_{\sigma(i)} + \cdots + \ell_{\sigma^{c-1}(i)}.$$

Then one can easily check that

$$\nu^{m_i} P_i \simeq P_i[n(m_i - c)]$$

for any $1 \le i \le a$.

(a) By Proposition 4.3(a), we only have to check $m_1 = \cdots = m_a$. Since Λ is ring-indecomposable, it is enough to check that $m_i = m_j$ holds if $\operatorname{Hom}_\Lambda(P_i, P_j) \neq 0$. In this case, we have

$$0 \neq \operatorname{Hom}_{\mathcal{D}}(\nu^{m_i m_j} P_i, \nu^{m_i m_j} P_j) = \operatorname{Hom}_{\mathcal{D}}(P_i[n(m_i - c)m_j], P_j[n(m_j - c)m_i]) = \operatorname{Hom}_{\mathcal{D}}(P_i, P_j[nc(m_j - m_i)]).$$

Thus we have $nc(m_j - m_i) = 0$ and $m_i = m_j$.

(b) We have shown that Λ is twisted $\frac{n(m-c)}{m}$-CY for $m := m_1 = \cdots = m_a$. Since

$$\frac{c}{m} = \frac{ac}{m_1 + \cdots + m_a} = \frac{a}{\ell_1 + \cdots + \ell_a} = \frac{a}{b},$$

we have the assertion. □

4.2. **Proof of Theorem 1.3.** (a)⇒(b) By Lemma 4.4(b), we have

$$\nu_n^{\ell - 1}(D\Lambda) \simeq \tau_n^{\ell - 1}(D\Lambda) \simeq \Lambda.$$

Thus we have

$$\nu^\ell \Lambda \simeq \nu^{\ell - 1}(D\Lambda) \simeq \Lambda[n(\ell - 1)].$$

The assertion follows from Proposition 4.3(a).

(b)⇒(a) We need the following results [IO2, 3.1] and [I4, 5.4(a)].

Proposition 4.5. *Let Λ be a finite dimensional K-algebra with* $\operatorname{gl.dim} \Lambda \le n$.

(a) *Let $\mathcal{U} := \operatorname{add}\{\nu_n^j \Lambda \mid j \in \mathbf{Z}\} \subset \mathcal{D}$. Then Λ is n-representation-finite if and only if $D\Lambda \in \mathcal{U}$.*

(b) *Let $(\mathcal{D}^{\le 0}, \mathcal{D}^{\ge 0})$ be the standard t-structure of $\mathcal{D}$. Then $\nu_n \mathcal{D}^{\ge 0} \subset \mathcal{D}^{\ge 0}$ and $\nu_n^{-1} \mathcal{D}^{\le 0} \subset \mathcal{D}^{\le 0}$.*

Since $\nu^\ell \Lambda \simeq \Lambda[n(\ell - 1)]$, we have $\nu_n^{\ell - 1}(D\Lambda) \simeq \Lambda$, which implies $D\Lambda \in \mathcal{U}$. By Proposition 4.5(a), we have that Λ is n-representation-finite. On the other hand, we have

$$\nu_n^{-i} \Lambda \simeq \nu_n^{\ell - 1 - i}(D\Lambda) \in \mathcal{D}^{\le 0} \cap \mathcal{D}^{\ge 0} = \operatorname{mod} \Lambda$$

for any $0 \le i < \ell$ by Proposition 4.5(b). By Lemma 4.4(a), we have $\tau_n^{\ell - 1}(D\Lambda) \simeq \nu_n^{\ell - 1}(D\Lambda) \simeq \Lambda$. Thus Λ is ℓ-homogeneous. □

4.3. **Proof of Proposition 1.4 and Corollary 1.5.** We shall prove Proposition 1.4.

Since K is a perfect field, $\Lambda_1 \otimes \cdots \otimes \Lambda_k$ has finite global dimension. We prove the assertion for twisted fractionally CY property. One can show the assertion for fractionally CY property similarly by using Proposition 4.3(b) instead of (a).

By Proposition 4.3(a), we have an isomorphism

$$(D\Lambda_i)^{\overset{\mathbf{L}}{\otimes}_{\Lambda_i} \ell} \simeq \Lambda_i[n_i(\ell-1)]$$

in $\mathcal{D}^{\mathrm{b}}(\operatorname{mod}\Lambda_i)$. Thus we have

$$\begin{aligned} &(D(\Lambda_1 \otimes \cdots \otimes \Lambda_k))^{\overset{\mathbf{L}}{\otimes}_{\Lambda_1 \otimes \cdots \otimes \Lambda_k} \ell} \simeq (D\Lambda_1)^{\overset{\mathbf{L}}{\otimes}_{\Lambda_1} \ell} \otimes \cdots \otimes (D\Lambda_k)^{\overset{\mathbf{L}}{\otimes}_{\Lambda_k} \ell} \\ \simeq \quad & \Lambda_1\left[\frac{\ell m_1}{\ell_1}\right] \otimes \cdots \otimes \Lambda_k\left[\frac{\ell m_k}{\ell_k}\right] \simeq (\Lambda_1 \otimes \cdots \otimes \Lambda_k)\left[\ell\left(\frac{m_1}{\ell_1} + \cdots \frac{m_k}{\ell_k}\right)\right] \end{aligned}$$

in $\mathcal{D}^{\mathrm{b}}(\operatorname{mod}\Lambda_1 \otimes \cdots \otimes \Lambda_k)$. By Proposition 4.3(a), we have the assertion. □

We shall prove Corollary 1.5.

First of all, note that $\Lambda_1 \otimes \cdots \otimes \Lambda_k$ has global dimension at most $n_1 + \cdots + n_k$ since K is a perfect field. By Theorem 1.3(a)⇒(b), we have that Λ_i is twisted $\frac{n_i(\ell-1)}{\ell}$-CY. By Proposition 1.4, we have that $\Lambda_1 \otimes \cdots \otimes \Lambda_k$ is twisted $\frac{(n_1+\cdots+n_k)(\ell-1)}{\ell}$-CY. By Theorem 1.3(b)⇒(a), we have that $\Lambda_1 \otimes \cdots \otimes \Lambda_k$ is ℓ-homogeneous $(n_1 + \cdots + n_k)$-representation-finite.

The second statement is clear from Proposition 0.2 since we have

$$\tau_{n_1+\cdots+n_k}(X_1 \otimes \cdots \otimes X_k) = (\tau_{n_1}X_1) \otimes \cdots \otimes (\tau_{n_k}X_k)$$

for any $X_i \in \operatorname{mod}\Lambda_i$. □

We end this section by giving another proof of Corollary 1.5.

Since Λ_i is ℓ-homogeneous n_i-representation-finite, we have $D\Lambda_i \simeq \nu_{n_i}^{-\ell}\Lambda_i$ for any i. Thus we have

$$D(\Lambda_1 \otimes \cdots \otimes \Lambda_k) \simeq D\Lambda_1 \otimes \cdots \otimes D\Lambda_k \simeq \nu_{n_1}^{-\ell}\Lambda_1 \otimes \cdots \otimes \nu_{n_k}^{-\ell}\Lambda_k \simeq \nu_{n_1+\cdots+n_k}^{-\ell}(\Lambda_1 \otimes \cdots \otimes \Lambda_k).$$

Thus $\Lambda_1 \otimes \cdots \otimes \Lambda_k$ is $(n_1 + \cdots + n_k)$-representation-finite by Proposition 4.5(a). □

4.4. **Proof of Proposition 1.7.** We have $M = \bigoplus_{i=0}^{\ell-1} \tau_n^{-i}\Lambda$ by Proposition 0.2. We have

$$\operatorname{Hom}_\Lambda(\tau_n^{-i}\Lambda, \tau_n^{-j}\Lambda) \overset{\text{Prop. } 0.2}{\simeq} \begin{cases} \operatorname{Hom}_\Lambda(\tau_n^{-(i-j)}\Lambda, \Lambda) \overset{\text{Lem. } 4.4(\mathrm{b})}{=} 0 & i > j, \\ \operatorname{Hom}_\Lambda(\Lambda, \tau_n^{-(j-i)}\Lambda) \simeq \tau_n^{j-i}\Lambda & i \leq j. \end{cases}$$

Since we have $\tau_n^{-i}\Lambda \simeq \nu_n^{-i}\Lambda \simeq T^{\otimes_\Lambda i}$ for $0 \leq i < \ell$ by Lemma 4.4(b) and [IO1, 2.12], we have the assertion. □

4.5. **Proof of results in Section 2.** Our Proposition 2.1 is an immediate consequence of the following observation.

Lemma 4.6. *Assume* $\operatorname{Hom}_\Lambda(P_i, P_j) \neq 0$*, or equivalently* $\operatorname{Hom}_\Lambda(I_i, I_j) \neq 0$*. Then we have* $\ell_i \leq \ell_j$ *and* $\ell_{\sigma^{-1}(i)} \geq \ell_{\sigma^{-1}(j)}$.

Proof. If $\ell_i > \ell_j$, then by Theorem 0.2, we have

$$0 \neq \operatorname{Hom}_\Lambda(I_i, I_j) \simeq \operatorname{Hom}_\Lambda(\tau_n^{\ell_j-1}I_i, P_{\sigma(j)})$$

by applying $\tau_n^{\ell_j-1}$. By Lemma 4.4(b), the right hand side is zero, a contradiction. Thus we have $\ell_i \leq \ell_j$.

If $\ell_{\sigma^{-1}(i)} < \ell_{\sigma^{-1}(j)}$, then we have

$$0 \neq \operatorname{Hom}_\Lambda(P_i, P_j) \simeq \operatorname{Hom}_\Lambda(I_{\sigma^{-1}(i)}, \tau_n^{-(\ell_{\sigma^{-1}(i)}-1)}P_j)$$

by applying $\tau_n^{-(\ell_{\sigma^{-1}(i)}-1)}$. By Lemma 4.4(b), the right hand side is zero, a contradiction. □

Next we give a proof of Proposition 2.2.

We have

$$D\operatorname{Hom}_{\mathcal{D}}(\nu_n^{j}\,P_i,\Lambda)\simeq\operatorname{Hom}_{\mathcal{D}}(\Lambda,\nu\,\nu_n^{j}\,P_i)\simeq\operatorname{Hom}_{\mathcal{D}}(\Lambda,\nu_n^{j}\,I_i)\simeq\operatorname{Hom}_{\mathcal{D}}(\Lambda,\nu_n^{j-\ell_i}\,P_{\sigma(i)}). \tag{2}$$

Since we have functorial isomorphisms

$$\Pi\otimes_\Lambda - \simeq \bigoplus_{j\in\mathbf{Z}}\operatorname{Hom}_{\mathcal{D}}(\Lambda,\nu_n^{j}\,-)\quad:\quad\operatorname{mod}\Lambda\to\operatorname{mod}\Pi, \tag{3}$$

$$\operatorname{Hom}_{\Pi}(\Pi\otimes_\Lambda -,\Pi)\simeq\bigoplus_{j\in\mathbf{Z}}\operatorname{Hom}_{\mathcal{D}}(\nu_n^{j}\,-,\Lambda)\quad:\quad\operatorname{mod}\Lambda\to\operatorname{mod}\Pi^{\mathrm{op}} \tag{4}$$

on $\operatorname{add}\Lambda$, we have

$$\psi^*(\Pi\otimes_\Lambda P_i)\overset{(4)}{\simeq}\bigoplus_{j\in\mathbf{Z}}D\operatorname{Hom}_{\mathcal{D}}(\nu_n^{j}\,P_i,\Lambda)\overset{(2)}{\simeq}\bigoplus_{j\in\mathbf{Z}}\operatorname{Hom}_{\mathcal{D}}(\Lambda,\nu_n^{j-\ell_i}\,P_{\sigma(i)})\overset{(3)}{\simeq}\Pi\otimes_\Lambda P_{\sigma(i)}.$$

□

Now we give a proof of Proposition 2.4.

We always identify $\phi^*\Lambda$ with Λ by the isomorphism $\phi^*\Lambda=\Lambda_\phi\otimes_\Lambda\Lambda\ni\lambda\otimes\gamma\mapsto\lambda\cdot\phi\gamma\in\Lambda$.

(i) First we define a K-linear automorphism $\widetilde{\phi}$ of Π by using the action of the autofunctor ϕ^* of $\mathcal{D}$ on $\operatorname{Hom}_{\mathcal{D}}(\Lambda,\nu_n^{-i}\,\Lambda)$.

By Lemma 4.4(c), we have an isomorphism $\rho:\phi^*\,\nu_n^{-1}\to\nu_n^{-1}\,\phi^*$ of autofunctors of $\mathcal{D}$. For any $i\geq 0$, we define an isomorphism $\rho^i:\phi^*\,\nu_n^{-i}\to\nu_n^{-i}\,\phi^*$ of autofunctors of $\mathcal{D}$ by $\rho^0=\mathrm{id}_{\phi^*}$ and

$$\rho^i:=(\phi^*\,\nu_n^{-i}\xrightarrow{\rho_{\nu_n^{1-i}}}\nu_n^{-1}\,\phi^*\,\nu_n^{1-i}\xrightarrow{\nu_n^{-1}\,\rho_{\nu_n^{2-i}}}\cdots\xrightarrow{\nu_n^{2-i}\,\rho_{\nu_n^{-1}}}\nu_n^{1-i}\,\phi^*\,\nu_n^{-1}\xrightarrow{\nu_n^{1-i}\,\rho}\nu_n^{-i}\,\phi^*).$$

Clearly we have

$$\rho^{i+j}=(\phi^*\,\nu_n^{-i-j}\xrightarrow{\rho^i_{\nu_n^{-j}}}\nu_n^{-i}\,\phi^*\,\nu_n^{-j}\xrightarrow{\nu_n^{-i}\,\rho^j}\nu_n^{-i-j}\,\phi^*) \tag{5}$$

for any $i,j\geq 0$. For any $f\in\operatorname{Hom}_{\mathcal{D}}(\Lambda,\nu_n^{-i}\,\Lambda)$, define $\widetilde{\phi}f\in\operatorname{Hom}_{\mathcal{D}}(\Lambda,\nu_n^{-i}\,\Lambda)$ by

$$\widetilde{\phi}f:=(\Lambda=\phi^*\Lambda\xrightarrow{\phi^*f}\phi^*\,\nu_n^{-i}\,\Lambda\xrightarrow{\rho^i_\Lambda}\nu_n^{-i}\,\phi^*\Lambda=\nu_n^{-i}\,\Lambda).$$

Clearly the restriction of $\widetilde{\phi}$ to Λ coincides with ϕ.

(ii) Next we show that $\widetilde{\phi}$ is a K-algebra automorphism of Π.

Fix $f\in\operatorname{Hom}_{\mathcal{D}}(\Lambda,\nu_n^{-i}\,\Lambda)$ and $g\in\operatorname{Hom}_{\mathcal{D}}(\Lambda,\nu_n^{-j}\,\Lambda)$. Since ρ^i is an isomorphism of functors, we have a commutative diagram

$$\begin{tikzcd}
\phi^*\,\nu_n^{-i}\,\Lambda \arrow[r, "\phi^*\,\nu_n^{-i}\,g"] \arrow[d, "\rho^i_\Lambda"] & \phi^*\,\nu_n^{-i-j}\,\Lambda \arrow[d, "\rho^i_{\nu_n^{-j}\,\Lambda}"] \\
\nu_n^{-i}\,\phi^*\Lambda \arrow[r, "\nu_n^{-i}\,\phi^*g"] & \nu_n^{-i-j}\,\phi^*\Lambda.
\end{tikzcd} \tag{6}$$

Now we have

$$\begin{aligned}
\widetilde{\phi}f\cdot\widetilde{\phi}g &= (\Lambda \xrightarrow{\widetilde{\phi}f} \nu_n^{-i}\,\Lambda \xrightarrow{\nu_n^{-i}(\widetilde{\phi}g)} \nu_n^{-i-j}\,\Lambda)\\
&= (\Lambda=\phi^*\Lambda \xrightarrow{\phi^* f} \phi^*\,\nu_n^{-i}\,\Lambda \xrightarrow{\rho^i_\Lambda} \nu_n^{-i}\,\phi^*\Lambda \xrightarrow{\nu_n^{-i}\,\phi^* g} \nu_n^{-i}\,\phi^*\,\nu_n^{-j}\,\Lambda =\\
&\quad\ \nu_n^{-i}\,\phi^*\,\nu_n^{-j}\,\Lambda \xrightarrow{\nu_n^{-i}\,\rho^j_\Lambda} \nu_n^{-i-j}\,\phi^*\Lambda=\nu_n^{-i-j}\,\Lambda)\\
&\overset{(6)}{=} (\Lambda=\phi^*\Lambda \xrightarrow{\phi^* f} \phi^*\,\nu_n^{-i}\,\Lambda \xrightarrow{\phi^*\,\nu_n^{-i}\,g} \phi^*\,\nu_n^{-i-j}\,\Lambda \xrightarrow{\rho^i_{\nu_n^{-j}\,\Lambda}} \nu_n^{-i}\,\phi^*\,\nu_n^{-j}\,\Lambda =\\
&\quad\ \nu_n^{-i}\,\phi^*\,\nu_n^{-j}\,\Lambda \xrightarrow{\nu_n^{-i}\,\rho^j_\Lambda} \nu_n^{-i-j}\,\phi^*\Lambda=\nu_n^{-i-j}\,\Lambda)\\
&\overset{(5)}{=} (\Lambda=\phi^*\Lambda \xrightarrow{\phi^*(f\,\nu_n^{-i}\,g)} \phi^*\,\nu_n^{-i-j}\,\Lambda \xrightarrow{\rho^{i+j}_\Lambda} \nu_n^{-i-j}\,\phi^*\Lambda=\nu_n^{-i-j}\,\Lambda)\\
&= \widetilde{\phi}(f\,\nu_n^{-i}\,g).
\end{aligned}$$

(iii) Since $\nu^\ell \simeq [n(\ell-1)]\circ\phi^*$, we have

$$\nu\circ\nu_n^{\ell-1}\simeq\phi^*. \tag{7}$$

By Serre duality, we have isomorphisms

$$D\operatorname{Hom}_{\mathcal{D}}(\Lambda,\nu_n^{-i}\,\Lambda)\simeq\operatorname{Hom}_{\mathcal{D}}(\nu_n^{-i}\,\Lambda,\nu\,\Lambda)\simeq\operatorname{Hom}_{\mathcal{D}}(\Lambda,\nu\,\nu_n^{i}\,\Lambda)\overset{(7)}{\simeq}\operatorname{Hom}_{\mathcal{D}}(\Lambda,\phi^*\,\nu_n^{i+1-\ell}\,\Lambda). \tag{8}$$

Thus we have an isomorphism

$$D\Pi=\bigoplus_{i\in\mathbf{Z}}D\operatorname{Hom}_{\mathcal{D}}(\Lambda,\nu_n^{-i}\,\Lambda)\overset{(8)}{\simeq}\bigoplus_{i\in\mathbf{Z}}\operatorname{Hom}_{\mathcal{D}}(\Lambda,\phi^*\,\nu_n^{-i}\,\Lambda)=:N \tag{9}$$

of $\Pi\otimes\Pi^{\mathrm{op}}$-modules, where Π acts on N from the left in a usual way, and from the right by

$$f\cdot g=(\Lambda\xrightarrow{f}\phi^*\,\nu_n^{-i}\,\Lambda\xrightarrow{\phi^*\,\nu_n^{-i}\,g}\phi^*\,\nu_n^{-i-j}\,\Lambda) \tag{10}$$

for any $f\in\operatorname{Hom}_{\mathcal{D}}(\Lambda,\phi^*\,\nu_n^{-i}\,\Lambda)\subset N$ and $g\in\operatorname{Hom}_{\mathcal{D}}(\Lambda,\nu_n^{-j}\,\Lambda)\subset\Pi$. Let

$$\gamma: N=\bigoplus_{i\in\mathbf{Z}}\operatorname{Hom}_{\mathcal{D}}(\Lambda,\phi^*\,\nu_n^{-i}\,\Lambda)\to\Pi_{\widetilde{\phi}}=\bigoplus_{i\in\mathbf{Z}}\operatorname{Hom}_{\mathcal{D}}(\Lambda,\nu_n^{-i}\,\Lambda), \tag{11}$$

be a K-linear isomorphism given by

$$\gamma f:=(\Lambda\xrightarrow{f}\phi^*\,\nu_n^{-i}\,\Lambda\xrightarrow{\rho^i_\Lambda}\nu_n^{-i}\,\phi^*\Lambda=\nu_n^{-i}\,\Lambda)$$

for any $f\in\operatorname{Hom}_{\mathcal{D}}(\Lambda,\phi^*\,\nu_n^{-i}\,\Lambda)$. Then γ is an isomorphism of $\Pi\otimes\Pi^{\mathrm{op}}$-modules by the following equalities for any $f\in\operatorname{Hom}_{\mathcal{D}}(\Lambda,\phi^*\,\nu_n^{-i}\,\Lambda)\subset N$ and $g\in\operatorname{Hom}_{\mathcal{D}}(\Lambda,\nu_n^{-j}\,\Lambda)\subset\Pi$.

$$\begin{aligned}
\gamma f\cdot\widetilde{\phi}g &= (\Lambda\xrightarrow{f}\phi^*\,\nu_n^{-i}\,\Lambda\xrightarrow{\rho^i_\Lambda}\nu_n^{-i}\,\phi^*\Lambda\xrightarrow{\nu_n^{-i}\,\phi^* g}\nu_n^{-i}\,\phi^*\,\nu_n^{-j}\,\Lambda\xrightarrow{\nu_n^{-i}\,\rho^j_\Lambda}\nu_n^{-i-j}\,\phi^*\Lambda=\nu_n^{-i-j}\,\Lambda)\\
&\overset{(6)}{=} (\Lambda\xrightarrow{f}\phi^*\,\nu_n^{-i}\,\Lambda\xrightarrow{\phi^*\,\nu_n^{-i}\,g}\phi^*\,\nu_n^{-i-j}\,\Lambda\xrightarrow{\rho^i_{\nu_n^{-j}\,\Lambda}}\nu_n^{-i}\,\phi^*\,\nu_n^{-j}\,\Lambda\xrightarrow{\nu_n^{-i}\,\rho^j_\Lambda}\nu_n^{-i-j}\,\phi^*\Lambda=\nu_n^{-i-j}\,\Lambda)\\
&\overset{(5)}{=} (\Lambda\xrightarrow{f}\phi^*\,\nu_n^{-i}\,\Lambda\xrightarrow{\phi^*\,\nu_n^{-i}\,g}\phi^*\,\nu_n^{-i-j}\,\Lambda\xrightarrow{\rho^{i+j}_\Lambda}\nu_n^{-i-j}\,\phi^*\Lambda=\nu_n^{-i-j}\,\Lambda)\\
&\overset{(10)}{=} \gamma(f\cdot g).
\end{aligned}$$

Combining (9) and (11), we have an isomorphism $D\Pi\simeq\Pi_{\widetilde{\phi}}$ of $\Pi\otimes\Pi^{\mathrm{op}}$-modules. Thus we have that $\widetilde{\phi}$ is a Nakayama automorphism of Π. □

Finally we give a proof of Theorem 2.3.

Assume that Λ is ℓ-homogeneous. By Proposition 2.4, the K-automorphism ϕ of Λ extends to a Nakayama automorphism $\widetilde{\phi}$ of Π, which satisfies $\widetilde{\phi}(I)=I$. Since $\widetilde{\phi}$ and ψ coincide in the outer automorphism group of Π, we have $\psi(I)=I$.

Assume $\psi(I)=I$. Then ψ induces a K-algebra automorphism $\psi|_\Lambda$ of Λ. By Proposition 2.2, we have $\psi|_\Lambda^*(P_i)\simeq P_{\sigma(i)}$ and $\psi|_\Lambda^*(I_i)\simeq I_{\sigma(i)}$ by Lemma 4.4(c). Since we have $\psi|_\Lambda^*(\tau_n^j I_i)=\tau_n^j\psi|_\Lambda^*(I_i)=\tau_n^j I_{\sigma(i)}$ for any j by Lemma 4.4(c), we have $\ell_i=\ell_{\sigma(i)}$. By Proposition 2.1, we have that Λ is homogeneous. □

References

[Am] C. Amiot, *Cluster categories for algebras of global dimension 2 and quivers with potential*, arXiv:0805.1035, to appear in Ann. Inst. Fourier.

[Au] M. Auslander, *Representation dimension of Artin algebras*, Lecture notes, Queen Mary College, London, 1971.

[Ba] *Spectral Methods in Representation Theory of Algebras and Applications to the Study of Rings of Singularities*, September 7-12, 2008, Banff International Research Station, Canada.

[Bi] *The ADE chain*, October 31-November 1, 2008, Universitaet Bielefeld, Germany.

[BK] A. I. Bondal, M. M. Kapranov, *Representable functors, Serre functors, and reconstructions*, Izv. Akad. Nauk SSSR Ser. Mat. 53 (1989), no. 6, 1183–1205, 1337; translation in Math. USSR-Izv. 35 (1990), no. 3, 519–541

[BIRS] A. Buan, O. Iyama, I. Reiten, J. Scott, *Cluster structure for 2-Calabi-Yau categories and unipotent groups*, Compos. Math. 145 (2009), no. 4, 1035–1079.

[BMRRT] A. Buan, R. Marsh, I. Reiten, M. Reineke, G. Todorov, *Tilting theory and cluster combinatorics*, Adv. in Math. 204 (2006), no. 2, 572–618.

[D] J. A. de la Pena, lectures at conferences [Ba, P].

[DL] H. Lenzing, J. A. de la Pena, *Extended canonical algebras and Fuchsian singularities*, arXiv:math.RT/0611532.

[EH] K. Erdmann, T. Holm, *Maximal n-orthogonal modules for selfinjective algebras*, Proc. Amer. Math. Soc. 136 (2008) no. 9, 3069–3078.

[G] P. Gabriel, *Auslander-Reiten sequences and representation-finite algebras*, Representation theory, I (Proc. Workshop, Carleton Univ., Ottawa, Ont., 1979), pp. 1–71, Lecture Notes in Math., 831, Springer, Berlin, 1980.

[GLS1] C. Geiss, B. Leclerc, J. Schröer, *Rigid modules over preprojective algebras*, Invent. Math. 165 (2006), no. 3, 589–632.

[GLS2] C. Geiss, B. Leclerc, J. Schröer, *Cluster algebra structures and semicanoncial bases for unipotent groups*, arXiv:math/0703039.

[H] D. Happel, *Triangulated categories in the representation theory of finite-dimensional algebras*, London Mathematical Society Lecture Note Series, 119. Cambridge University Press, Cambridge, 1988.

[HZ1] Z. Huang, X. Zhang, *The existence of maximal n-orthogonal subcategories*, J. Algebra 321 (2009), no. 10, 2829–2842.

[HZ2] Z. Huang, X. Zhang, *Higher Auslander algebras admitting trivial maximal orthogonal subcategories*, arXiv:0903.0761.

[HZ3] Z. Huang, X. Zhang, *Trivial maximal 1-orthognal subcategories for Auslander's 1-Gorenstein algebras*, arXiv:0903.0762.

[IIKNS] R. Inoue, O. Iyama, A. Kuniba, T. Nakanishi, J. Suzuki, *Periodicities of T-systems and Y-systems*, to appear in Nagoya Math. J., arXiv:0812.0667.

[I1] O. Iyama, *Higher-dimensional Auslander-Reiten theory on maximal orthogonal subcategories*, Adv. Math. 210 (2007), no. 1, 22–50.

[I2] O. Iyama, *Auslander correspondence*, Adv. Math. 210 (2007), no. 1, 51–82.

[I3] O. Iyama, *Auslander-Reiten theory revisited*, Trends in Representation Theory of Algebras and Related Topics, 349–398, European Mathematical Society, 2008.

[I4] O. Iyama, *Cluster tilting for higher Auslander algebras*, arXiv:0809.4897.

[IKM] O. Iyama, K. Kato, J. Miyachi, *Recollement of homotopy categories and Cohen-Macaulay modules*, arXiv:0911.0172.

[IO1] O. Iyama, S. Oppermann, *n-representation-finite algebras and n-APR tilting*, to appear in Trans. Amer. Math. Soc., arXiv:0909.0593.

[IO2] O. Iyama, S. Oppermann, *Stable categories over higher preprojective algebras*, arXiv:0912.3412.

[IR] O. Iyama, I. Reiten, *Fomin-Zelevinsky mutation and tilting modules over Calabi-Yau algebras*, Amer. J. Math. 130 (2008), no. 4, 1087–1149.

[KST] H. Kajiura, K. Saito, A. Takahashi, *Matrix factorization and representations of quivers. II. Type ADE case*, Adv. Math. 211 (2007), no. 1, 327–362.

[Ke1] B. Keller, *Calabi-Yau triangulated categories*, Trends in Representation Theory of Algebras and Related Topics, 467–489, European Mathematical Society, 2008.

[Ke2] B. Keller, *Cluster algebras, quiver representations and triangulated categories*, arXiv:0807.1960.

[Ke3] B. Keller, *Deformed Calabi-Yau completions*, arXiv:0908.3499.

[KR] B. Keller, I. Reiten, *Cluster-tilted algebras are Gorenstein and stably Calabi-Yau*, Adv. Math. 211 (2007), 123-151.

[Ko] M. Kontsevich, *Triangulated categories and geometry*, Course at the École Normale Supérieure, Paris, 1998.

[Lada] M. Lada, *Relative homology and maximal l-orthogonal modules*, J. Algebra 321 (2009), no. 10, 2798–2811.

[Ladk] S. Ladkani, lectures at conferences [Ba, Bi, P].

[Le] H. Lenzing, lectures at conferences [Ba, Bi, T].

[LM] H. Lenzing, H. Meltzer, *Sheaves on a weighted projective line of genus one, and representations of a tubular algebra*, Representations of algebras (Ottawa, ON, 1992), 313–337, CMS Conf. Proc., 14, Amer. Math. Soc., Providence, RI, 1993.

[Mi] H. Minamoto *Ampleness of two-sided tilting complexes*, preprint.

[MY] J. Miyachi, A. Yekutieli, *Derived Picard groups of finite-dimensional hereditary algebras*, Compositio Math. 129 (2001), no. 3, 341–368.

[P] *Workshop on Triangulated Categories and Singularities*, May 25-30, 2009, Universitaet Paderborn, Germany.

[R] C. M. Ringel, *Tame algebras and integral quadratic forms*, Lecture Notes in Mathematics, 1099. Springer-Verlag, Berlin, 1984.

[T] *Homological and geometric methods in algebra*, August 10-14, 2009, Norwegian University of Technology and Science, Norway.

[Y] K. Yanagawa, lectures at Nagoya University, 2009.

O. Iyama: Graduate School of Mathematics, Nagoya University, Chikusa-ku, Nagoya, 464-8602 Japan
E-mail address: `iyama@math.nagoya-u.ac.jp`

M. Herschend: Graduate School of Mathematics, Nagoya University, Chikusa-ku, Nagoya, 464-8602 Japan
E-mail address: `martin.herschend@gmail.com`